

[原著論文]

英文原著論文 (直近のものから順に記載。 *: corresponding author.)

1. Hori H*, Fukushima H, Nagayoshi T, Ishikawa R, Zhuo M, Yoshida F, Kunugi H, Okamoto K, Kim Y*, Kida S*. Fear memory regulation by the cAMP signaling pathway as an index of reexperiencing symptoms in posttraumatic stress disorder. *Mol Psychiatry* (online ahead of print).
2. Hori H*, Yoshida F, Ishida I, Matsuo J, Ogawa S, Hattori K, Kim Y, Kunugi H*. Blood mRNA expression levels of glucocorticoid receptors and FKBP5 are associated with depressive disorder and altered HPA axis. *J Affect Disord*. 2024; 349: 244-253.
3. Kawanishi H*, Hori H*, Yoshida F, Itoh M, Lin M, Niwa M, Narita M, Otsuka T, Ino K, Imai R, Fukudo S, Kamo T, Kunugi H, Kim Y. Suicidality in civilian women with PTSD: Possible link to childhood maltreatment, proinflammatory molecules, and their genetic variations. *Brain Behav Immun Health*. 2023; 30: 100650.
4. Hakamata Y*, Hori H, Mizukami S, Izawa S, Yoshida F, Moriguchi Y, Hanakawa T, Inoue Y, Tagaya H. Blunted diurnal interleukin-6 rhythm is associated with amygdala emotional hyporeactivity and depression: a modulating role of gene-stressor interactions. *Front Psychiatry*. 2023; 14: 1196235.
5. Yamada R, Fujii T, Hattori K, Hori H, Matsumura R, Kurashimo T, Ishihara N, Yoshida S, Sumiyoshi T, Kunugi H*. Discrepancy between clinician-rated and self-reported depression severity is associated with adverse childhood experience, autistic-like traits, and coping styles in mood disorders. *Clin Psychopharmacol Neurosci*. 2023; 21: 296-303.
6. Shiwaku H*, Katayama S, Gao M, Kondo K, Nakano Y, Motokawa Y, Toyoda S, Yoshida F, Hori H, Kubota T, Ishikawa K, Kunugi H, Ikegaya Y, Okazawa H, Takahashi H. Analyzing schizophrenia-related phenotypes in mice caused by autoantibodies against NRXN1 α in schizophrenia. *Brain Behav Immun*. 2023; 111: 32-45.
7. Kakehi R, Hori H*, Yoshida F, Itoh M, Lin M, Niwa M, Narita M, Ino K, Imai R, Sasayama D, Kamo T, Kunugi H, Kim Y. Hypothalamic-pituitary-adrenal axis and renin-angiotensin-aldosterone system in adulthood PTSD and childhood maltreatment history. *Front Psychiatry*. 2023; 13: 967779.
8. Narita Z, Nozaki S, Shikimoto R, Hori H, Kim Y, Mimura M, Tsugane S, Sawada N*. Association between vegetable, fruit, and flavonoid-rich fruit consumption in midlife and major depressive disorder in later life: the JPHC Saku Mental Health Study. *Transl Psychiatry*. 2022; 12: 412.
9. Nakajima M*, Hori H, Itoh M, Lin M, Kawanishi H, Narita M, Kim Y. Validation of childhood trauma questionnaire-short form in Japanese clinical and nonclinical adults. *Psychiatry Res Commun*. 2022; 2: 100065.
10. Niwa M*, Kato T, Narita-Ohtaki R, Otomo R, Suga Y, Sugawara M, Narita Z, Hori H, Kamo T, Kim Y. Skills Training in Affective and Interpersonal Regulation Narrative Therapy for women with ICD-11 complex PTSD related to childhood abuse in Japan: a pilot study. *Eur J Psychotraumatol*. 2022; 13: 2080933.
11. Ali AM*, Hori H, Kim Y, Kunugi H. The Depression Anxiety Stress Scale 8-Items expresses robust psychometric properties as an ideal shorter version of the Depression Anxiety Stress Scale 21 among

- healthy respondents from three continents. *Front Psychol.* 2022; 13: 799769.
12. Hori H*, Izawa S, Yoshida F, Kunugi H, Kim Y, Mizukami S, Inoue Y, Tagaya H, Hakamata Y*. Association of childhood maltreatment history with salivary interleukin-6 diurnal patterns and C-reactive protein in healthy adults. *Brain Behav Immun.* 2022; 101: 377-382.
 13. Ali AM*, Hori H, Kim Y, Kunugi H. Predictors of nutritional status, depression, internet addiction, Facebook addiction, and tobacco smoking among women with eating disorders in Spain. *Front Psychiatry.* 2021; 12: 735109.
 14. Hakamata Y*, Mizukami S, Izawa S, Okamura H, Mihara K, Marusak H, Moriguchi Y, Hori H, Hanakawa T, Inoue Y, Tagaya H. Implicit and explicit emotional memory recall in anxiety and depression: Role of basolateral amygdala and cortisol-norepinephrine interaction. *Psychoneuroendocrinology.* 2022; 136: 105598.
 15. Ali AM*, Alkhamees AA, Hori H, Kim Y, Kunugi H. The Depression Anxiety Stress Scale 21: Development and validation of the Depression Anxiety Stress Scale 8-Item in psychiatric patients and the general public for easier mental health measurement in a post COVID-19 world. *Int J Environ Res Public Health.* 2021; 18: 10142.
 16. Saito K*, Hattori K, Hidese S, Sasayama D, Miyakawa T, Matsumura R, Tatsumi M, Yokota Y, Ota M, Hori H, Kunugi H*. Profiling of cerebrospinal fluid lipids and their relationship with plasma lipids in healthy humans. *Metabolites.* 2021; 11: 268.
 17. Hakamata Y*, Mizukami S, Izawa S, Moriguchi Y, Hori H, Matsumoto N, Hanakawa T, Inoue Y, Tagaya H. Childhood trauma affects autobiographical memory deficits through basal cortisol and prefrontal-extrastriate functional connectivity. *Psychoneuroendocrinology.* 2021; 127: 105172.
 18. Hamatani S, Hirano Y, Sugawara A, Isobe M, Kodama N, Yoshihara K, Moriguchi Y, Ando T, Endo Y, Takahashi J, Nohara N, Takamura T, Hori H, Noda T, Tose K, Watanabe K, Adachi H, Gondo M, Takakura S, Fukudo S, Shimizu E, Yoshiuchi K, Sato Y, Sekiguchi A*. Eating Disorder Neuroimaging Initiative (EDNI): a multicentre prospective cohort study protocol for elucidating the neural effects of cognitive-behavioural therapy for eating disorders. *BMJ Open.* 2021; 11: e042685.
 19. Hori H*, Itoh M, Lin M, Yoshida F, Niwa M, Hakamata Y, Matsui M, Kunugi H, Kim Y. Childhood maltreatment history and attention bias variability in healthy adult women: role of inflammation and the BDNF Val66Met genotype. *Transl Psychiatry.* 2021; 11: 122.
 20. Matsuo J, Hori H, Ishida I, Hiraishi M, Ota M, Hidese S, Yomogida Y, Kunugi H*. Performance on the Wechsler Adult Intelligence Scale (WAIS) in Japanese patients with bipolar and major depressive disorders in euthymic and depressed states. *Psychiatry Clin Neurosci.* 2021; 75: 128-137.
 21. Hori H, Itoh M, Matsui M, Kamo T, Saito T, Nishimatsu Y, Kito S, Kida S, Kim Y*. The efficacy of memantine in the treatment of civilian posttraumatic stress disorder: an open-label trial. *Eur J Psychotraumatol.* 2021; 12: 1859821.
 22. Otsuka T, Hori H*, Yoshida F, Itoh M, Lin M, Niwa M, Ino K, Imai R, Ogawa S, Matsui M, Kamo T, Kunugi H, Kim Y. Association of *CRP* genetic variation with symptomatology, cognitive function, and circulating proinflammatory markers in civilian women with PTSD. *J Affect Disord.* 2021; 279: 640-649.
 23. Nakayama M, Hori H*, Itoh M, Lin M, Niwa M, Ino K, Imai R, Ogawa S, Sekiguchi A, Matsui M, Kunugi H, Kim Y. Possible long-term effects of childhood maltreatment on cognitive function in adult women with posttraumatic stress disorder. *Front Psychiatry.* 2020; 11: 344.
 24. Hakamata Y*, Mizukami S, Izawa S, Moriguchi Y, Hori H, Kim Y, Hanakawa T, Inoue Y, Tagaya H. Basolateral amygdala connectivity with subgenual anterior cingulate cortex represents enhanced

- fear-related memory encoding in anxious humans. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2020; 5: 301-310.
25. Hori H*, Itoh M, Yoshida F, Lin M, Niwa M, Hakamata Y, Ino K, Imai R, Ogawa S, Matsui M, Kamo T, Kunugi H, Kim Y. The *BDNF* Val66Met polymorphism affects negative memory bias in civilian women with PTSD. *Sci Rep*. 2020; 10: 3151.
 26. Hori H*, Yoshida F, Itoh M, Lin M, Niwa M, Ino K, Imai R, Ogawa S, Matsui M, Kamo T, Kunugi H, Kim Y. Proinflammatory status-stratified blood transcriptome profiling of civilian women with PTSD. *Psychoneuroendocrinology*. 2020; 111: 104491.
 27. Tsuchimine S, Hattori K, Ota M, Hidese S, Teraishi T, Sasayama D, Hori H, Noda T, Yoshida S, Yoshida F, Kunugi H*. Reduced plasma orexin-A levels in patients with bipolar disorder. *Neuropsychiatr Dis Treat*. 2019; 15: 2221-2230.
 28. Koga N, Ogura J, Yoshida F, Hattori K, Hori H, Aizawa E, Ishida I, Kunugi H*. Altered polyunsaturated fatty acid levels in relation to proinflammatory cytokines, fatty acid desaturase genotype, and diet in bipolar disorder. *Transl Psychiatry*. 2019; 9: 208.
 29. Imai R, Hori H*, Itoh M, Lin M, Niwa M, Ino K, Ogawa S, Sekiguchi A, Kunugi H, Akechi T, Kamo T, Kim Y. Relationships of blood proinflammatory markers with psychological resilience and quality of life in civilian women with posttraumatic stress disorder. *Sci Rep*. 2019; 9: 17905.
 30. Hidese S, Ota M, Hori H, Matsuo J, Ishida I, Hiraishi M, Teraishi T, Hattori K, Kunugi H*. The relationship between the Wechsler Memory Scale-Revised scores and whole-brain structure in patients with schizophrenia and healthy individuals. *Cogn Neuropsychiatry*. 2019; 24: 80-91.
 31. Itoh M*, Hori H*, Lin M, Niwa M, Ino K, Imai R, Ogawa S, Matsui M, Kamo T, Kim Y. Memory bias and its association with memory function in women with posttraumatic stress disorder. *J Affect Disord*. 2019; 245: 461-467.
 32. Ota M*, Matsuo J, Ishida I, Takano H, Yokoi Y, Hori H, Yoshida S, Ashida K, Nakamura K, Takahashi T, Kunugi H. Effects of a medium-chain triglyceride-based ketogenic formula on cognitive function in patients with mild-to-moderate Alzheimer's disease. *Neurosci Lett*. 2019; 690: 232-236.
 33. Matsuo J, Ota M, Hidese S, Teraishi T, Hori H, Ishida I, Hiraishi M, Kunugi H*. Sensorimotor gating in depressed and euthymic patients with bipolar disorder: analysis on prepulse inhibition of acoustic startle response stratified by gender and state. *Front Psychiatry*. 2018; 9: 123.
 34. Hori H*, Nakamura S, Yoshida F, Teraishi T, Sasayama D, Ota M, Hattori K, Kim Y, Higuchi T, Kunugi H*. Integrated profiling of phenotype and blood transcriptome for stress vulnerability and depression. *J Psychiatr Res*. 2018; 104: 202-210.
 35. Ota M*, Matsuo J, Sato N, Teraishi T, Hori H, Hattori K, Kamio Y, Maikusa N, Matsuda H, Kunugi H. Relationship between autistic spectrum trait and regional cerebral blood flow in healthy male subjects. *Psychiatry Investig*. 2018; 15: 956-961.
 36. Narita-Ohtaki R, Hori H*, Itoh M, Lin M, Niwa M, Ino K, Imai R, Ogawa S, Sekiguchi A, Matsui M, Kunugi H, Kamo T, Kim Y. Cognitive function in Japanese women with posttraumatic stress disorder: Association with exercise habits. *J Affect Disord*. 2018; 236: 306-312.
 37. Imai R, Hori H*, Itoh M, Lin M, Niwa M, Ino K, Ogawa S, Ishida M, Sekiguchi A, Matsui M, Kunugi H, Akechi T, Kamo T, Kim Y. Inflammatory markers and their possible effects on cognitive function in women with posttraumatic stress disorder. *J Psychiatr Res*. 2018; 102: 192-200.
 38. Ogawa S, Koga N, Hattori K, Matsuo J, Ota M, Hori H, Sasayama D, Teraishi T, Ishida I, Yoshida F,

- Yoshida S, Noda T, Higuchi T, Kunugi H*. Plasma amino acid profile in major depressive disorder: Analyses in two independent case-control sample sets. *J Psychiatr Res.* 2018; 96: 23-32.
39. Ishii T, Hattori K, Miyakawa T, Watanabe K, Hidese S, Sasayama D, Ota M, Teraishi T, Hori H, Yoshida S, Nunomura A, Nakagome K, Kunugi H*. Increased cerebrospinal fluid complement C5 levels in major depressive disorder and schizophrenia. *Biochem Biophys Res Commun.* 2018; 497: 683-688.
 40. Teraishi T, Kajiwara M, Hori H, Sasayama D, Hidese S, Matsuo J, Ishida I, Kajiwara Y, Ozeki Y, Ota M, Hattori K, Higuchi T, Kunugi H*. 13C-phenylalanine breath test and serum biopterin in schizophrenia, bipolar disorder and major depressive disorder. *J Psychiatr Res.* 2018; 99: 142-150.
 41. Hidese S, Hattori K, Sasayama D, Miyakawa T, Matsumura R, Yokota Y, Ishida I, Matsuo J, Noda T, Yoshida S, Teraishi T, Hori H, Ota M, Kunugi H*. Cerebrospinal fluid neural cell adhesion molecule levels and their correlation with clinical variables in patients with schizophrenia, bipolar disorder, and major depressive disorder. *Prog Neuropsychopharmacol Biol Psychiatry.* 2017; 76: 12-18.
 42. Mizuno M*, Yamaguchi S, Taneda A, Hori H, Aikawa A, Fujii C. Development of Japanese version of King's Stigma Scale and its short version: Psychometric properties of a self-stigma measure. *Psychiatry Clin Neurosci.* 2017; 71: 189-197.
 43. Matsuo J, Ota M, Hidese S, Hori H, Teraishi T, Ishida I, Hiraishi M, Kunugi H*. Sexually dimorphic deficits of prepulse inhibition in patients with major depressive disorder and their relationship to symptoms: A large single ethnicity study. *J Affect Disord.* 2017; 211: 75-82.
 44. Sasayama D, Hattori K, Ogawa S, Yokota Y, Matsumura R, Teraishi T, Hori H, Ota M, Yoshida S, Kunugi H*. Genome-wide quantitative trait loci mapping of the human cerebrospinal fluid proteome. *Hum Mol Genet.* 2017; 26: 44-51.
 45. Ota M*, Matsuo J, Sato N, Teraishi T, Hori H, Hattori K, Kamio Y, Kunugi H. Correlation of reduced social communicational and interactional skills with regional grey matter volumes in schizophrenia patients. *Acta Neuropsychiatr.* 2017; 29: 374-381.
 46. Ninomiya-Baba M, Matsuo J, Sasayama D, Hori H, Teraishi T, Ota M, Hattori K, Noda T, Ishida I, Shibata S, Kunugi H*. Association of body mass index-related single nucleotide polymorphisms with psychiatric disease and memory performance in a Japanese population. *Acta Neuropsychiatr.* 2017; 29: 299-308.
 47. Hori H*, Teraishi T, Nagashima A, Koga N, Ota M, Hattori K, Kim Y, Higuchi T, Kunugi H. A personality-based latent class typology of outpatients with major depressive disorder: Association with symptomatology, prescription pattern and social function. *J Affect Disord.* 2017; 217: 8-15.
 48. Hidese S, Hattori K, Sasayama D, Miyakawa T, Matsumura R, Yokota Y, Ishida I, Matsuo J, Noda T, Yoshida S, Teraishi T, Hori H, Ota M, Kunugi H*. Cerebrospinal fluid neural cell adhesion molecule levels and their correlation with clinical variables in patients with schizophrenia, bipolar disorder, and major depressive disorder. *Prog Neuropsychopharmacol Biol Psychiatry.* 2017; 76: 12-18.
 49. Matsuo J, Ota M, Hori H, Hidese S, Teraishi T, Ishida I, Hiraishi M, Kunugi H*. A large single ethnicity study of prepulse inhibition in schizophrenia: Separate analysis by sex focusing on effect of symptoms. *J Psychiatr Res.* 2016; 82: 155-162.
 50. Ota M*, Hori H, Sato N, Yoshida F, Hattori K, Teraishi T, Kunugi H. Effects of ankyrin 3 gene risk variants on brain structures in patients with bipolar disorder and healthy subjects. *Psychiatry Clin Neurosci.* 2016; 70: 498-506.
 51. Hori H*, Koga N, Hidese S, Nagashima A, Kim Y, Higuchi T, Kunugi H. 24-h activity rhythm and sleep in depressed outpatients. *J Psychiatr Res.* 2016; 77: 27-34.

52. [Hori H*](#), Sasayama D, Teraishi T, Yamamoto N, Nakamura S, Ota M, Hattori K, Kim Y, Higuchi T, Kunugi H. Blood-based gene expression signatures of medication-free outpatients with major depressive disorder: integrative genome-wide and candidate gene analyses. *Sci Rep.* 2016; 6: 18776.
53. Teraishi T, [Hori H](#), Sasayama D, Matsuo J, Ogawa S, Ota M, Hattori K, Kajiwara M, Higuchi T, Kunugi H*. 13C-tryptophan breath test detects increased catabolic turnover of tryptophan along the kynurenine pathway in patients with major depressive disorder. *Sci Rep.* 2015; 5: 15994.
54. Hattori K*, Ota M, Sasayama D, Yoshida S, Matsumura R, Miyakawa T, Yokota Y, Yamaguchi S, Noda T, Teraishi T, [Hori H](#), Higuchi T, Kohsaka S, Goto Y, Kunugi H. Increased cerebrospinal fluid fibrinogen in major depressive disorder. *Sci Rep.* 2015; 5: 11412.
55. Ota M*, Noda T, Sato N, Okazaki M, Ishikawa M, Hattori K, [Hori H](#), Sasayama D, Teraishi T, Sone D, Kunugi H. Effect of electroconvulsive therapy on gray matter volume in major depressive disorder. *J Affect Disord.* 2015; 186: 186-191.
56. Ota M*, Wakabayashi C, Sato N, [Hori H](#), Hattori K, Teraishi T, Ozawa H, Okubo T, Kunugi H. Effect of l-theanine on glutamatergic function in patients with schizophrenia. *Acta Neuropsychiatr.* 2015; 27: 291-296.
57. Ota M*, Noda T, Sato N, Hattori K, [Hori H](#), Sasayama D, Teraishi T, Nagashima A, Obu S, Higuchi T, Kunugi H. White matter abnormalities in major depressive disorder with melancholic and atypical features: A diffusion tensor imaging study. *Psychiatry Clin Neurosci.* 2015; 69: 360-368.
58. Sasayama D*, [Hori H](#), Nakamura S, Yamamoto N, Hattori K, Teraishi T, Ota M, Kunugi H. Increased protein and mRNA expression of resistin after dexamethasone administration. *Horm Metab Res.* 2015; 47: 433-438.
59. Matsuo J, Kamio Y, Takahashi H, Ota M, Teraishi T, [Hori H](#), Nagashima A, Takei R, Higuchi T, Motohashi N, Kunugi H*. Autistic-like traits in adult patients with mood disorders and schizophrenia. *PLoS One.* 2015; 10: e0122711.
60. Ogawa S, Hattori K, Sasayama D, Yokota Y, Matsumura R, Matsuo J, Ota M, [Hori H](#), Teraishi T, Yoshida S, Noda T, Ohashi Y, Sato H, Higuchi T, Motohashi N, Kunugi H*. Reduced cerebrospinal fluid ethanolamine concentration in major depressive disorder. *Sci Rep.* 2015; 5: 7796.
61. Teraishi T, [Hori H](#), Sasayama D, Matsuo J, Ogawa S, Ishida I, Nagashima A, Kinoshita Y, Ota M, Hattori K, Higuchi T, Kunugi H*. Personality in remitted major depressive disorder with single and recurrent episodes assessed with the Temperament and Character Inventory. *Psychiatry Clin Neurosci.* 2015; 69: 3-11.
62. Fujii T, Ota M, [Hori H](#), Hattori K, Teraishi T, Matsuo J, Kinoshita Y, Ishida I, Nagashima A, Kunugi H*. The common functional FKBP5 variant rs1360780 is associated with altered cognitive function in aged individuals. *Sci Rep.* 2014; 4: 6696.
63. Ogawa S, Fujii T, Koga N, [Hori H](#), Teraishi T, Hattori K, Noda T, Higuchi T, Motohashi N, Kunugi H*. Plasma L-tryptophan concentration in major depressive disorder: new data and meta-analysis. *J Clin Psychiatry.* 2014; 75: e906-15.
64. Teraishi T, [Hori H](#), Sasayama D, Matsuo J, Ogawa S, Ishida I, Nagashima A, Kinoshita Y, Ota M, Hattori K, Kunugi H*. Relationship between Lifetime Suicide Attempts and Schizotypal Traits in Patients with Schizophrenia. *PLoS One.* 2014; 9: e107739.
65. Ota M*, Sato N, Sakai K, Okazaki M, Maikusa N, Hattori K, [Hori H](#), Teraishi T, Shimoji K, Yamada K, Kunugi H. Altered coupling of regional cerebral blood flow and brain temperature in schizophrenia compared with bipolar disorder and healthy subjects. *J Cereb Blood Flow Metab.* 2014; 34: 1868-1872.

66. Fujii T, Ota M, Hori H, Hattori K, Teraishi T, Sasayama D, Higuchi T, Kunugi H*. Association between the common functional *FKBP5* variant (rs1360780) and brain structure in a non-clinical population. *J Psychiatr Res.* 2014; 58: 96-101.
67. Hori H*, Fujii T, Yamamoto N, Teraishi T, Ota M, Matsuo J, Kinoshita Y, Ishida I, Hattori K, Okazaki M, Arima K, Kunugi H. Temperament and character in remitted and symptomatic patients with schizophrenia: modulation by the *COMT* Val158Met genotype. *J Psychiatr Res.* 2014; 56: 82-89.
68. Fujii K, Ozeki Y*, Okayasu H, Takano Y, Shinozaki T, Hori H, Orui M, Horie M, Kunugi H, Shimoda K. QT is longer in drug-free patients with schizophrenia compared with age-matched healthy subjects. *PLoS One.* 2014; 9: e98555.
69. Ota M*, Noda T, Sato N, Hattori K, Teraishi T, Hori H, Nagashima A, Shimoji K, Higuchi T, Kunugi H. Characteristic distributions of regional cerebral blood flow changes in major depressive disorder patients: a pseudo-continuous arterial spin labeling (pCASL) study. *J Affect Disord.* 2014; 165: 59-63.
70. Hori H*, Yamamoto N, Teraishi T, Ota M, Fujii T, Sasayama D, Matsuo J, Kinoshita Y, Hattori K, Nagashima A, Ishida I, Koga N, Higuchi T, Kunugi H. Cognitive effects of the *ANK3* risk variants in patients with bipolar disorder and healthy individuals. *J Affect Disord.* 2014; 158: 90-96.
71. Ota M*, Ishikawa M, Sato N, Okazaki M, Maikusa N, Hori H, Hattori K, Teraishi T, Ito K, Kunugi H. Pseudo-continuous arterial spin labeling MRI study of schizophrenic patients. *Schizophr Res.* 2014; 154: 113-118.
72. Fujii T, Hori H, Ota M, Hattori K, Teraishi T, Sasayama D, Yamamoto N, Higuchi T, Kunugi H*. Effect of the common functional *FKBP5* variant (rs1360780) on the hypothalamic-pituitary-adrenal axis and peripheral blood gene expression. *Psychoneuroendocrinology.* 2014; 42: 89-97.
73. Richards M, Hori H*, Sartorius N, Kunugi H. Cross-cultural comparisons of attitudes toward schizophrenia amongst the general population and physicians: A series of web-based surveys in Japan and the United States. *Psychiatry Res.* 2014; 215: 300-307.
74. Ota M*, Wakabayashi C, Matsuo J, Kinoshita Y, Hori H, Hattori K, Sasayama D, Teraishi T, Obu S, Ozawa H, Kunugi H. Effect of L-theanine on sensorimotor gating in healthy human subjects. *Psychiatry Clin Neurosci.* 2014; 68: 337-343.
75. Sasayama D, Hori H, Yamamoto N, Nakamura S, Teraishi T, Tatsumi M, Hattori K, Ota M, Higuchi T, Kunugi H*. *ITIH3* polymorphism may confer susceptibility to psychiatric disorders by altering the expression levels of *GLT8D1*. *J Psychiatr Res.* 2014; 50: 79-83.
76. Sasayama D, Hori H, Teraishi T, Hattori K, Ota M, Matsuo J, Kinoshita Y, Okazaki M, Arima K, Amano N, Higuchi T, Kunugi H*. Benzodiazepines, benzodiazepine-like drugs, and typical antipsychotics impair manual dexterity in patients with schizophrenia. *J Psychiatr Res.* 2014; 49: 37-42.
77. Hori H*, Teraishi T, Ota M, Hattori K, Matsuo J, Kinoshita Y, Ishida I, Nagashima A, Koga N, Higuchi T, Kunugi H. Psychological coping in depressed outpatients: association with cortisol response to the combined dexamethasone/CRH test. *J Affect Disord.* 2014; 152-154: 441-447.
78. Hori H*, Teraishi T, Sasayama D, Matsuo J, Kinoshita Y, Ota M, Hattori K, Kunugi H. A latent profile analysis of schizotypy, temperament and character in a nonclinical population: association with neurocognition. *J Psychiatr Res.* 2014; 48: 56-64.
79. Ota M*, Hori H, Sato N, Sasayama D, Hattori K, Teraishi T, Obu S, Nakata Y, Kunugi H. Hypothalamic-pituitary-adrenal axis hyperactivity and brain differences in healthy women. *Neuropsychobiology.* 2013; 68: 205-211.
80. Teraishi T, Sasayama D, Hori H, Yamamoto N, Fujii T, Matsuo J, Nagashima A, Kinoshita Y, Hattori K,

- Ota M, Fujii S, Kunugi H*. Possible association between common variants of the phenylalanine hydroxylase (PAH) gene and memory performance in healthy adults. *Behav Brain Funct.* 2013; 9: 30.
81. Ota M*, Ishikawa M, Sato N, Hori H, Sasayama D, Hattori K, Teraishi T, Noda T, Obu S, Nakata Y, Higuchi T, Kunugi H. Discrimination between schizophrenia and major depressive disorder by magnetic resonance imaging of the female brain. *J Psychiatr Res.* 2013; 47: 1383-1388.
 82. Sasayama D, Hori H, Nakamura S, Miyata R, Teraishi T, Hattori K, Ota M, Yamamoto N, Higuchi T, Amano N, Kunugi H*. Identification of single nucleotide polymorphisms regulating peripheral blood mRNA expression with genome-wide significance: an eQTL study in the Japanese population. *PLoS One.* 2013; 8: e54967.
 83. Ota M*, Sato N, Matsuo J, Kinoshita Y, Kawamoto Y, Hori H, Teraishi T, Sasayama D, Hattori K, Obu S, Nakata Y, Kunugi H. Multimodal image analysis of sensorimotor gating in healthy women. *Brain Res.* 2013; 1499: 61-68.
 84. Sasayama D, Hattori K, Wakabayashi C, Teraishi T, Hori H, Ota M, Yoshida S, Arima K, Higuchi T, Amano N, Kunugi H*. Increased cerebrospinal fluid interleukin-6 levels in patients with schizophrenia and those with major depressive disorder. *J Psychiatr Res.* 2013; 47: 401-406.
 85. Hori H*, Matsuo J, Teraishi T, Sasayama D, Kawamoto Y, Kinoshita Y, Ota M, Hattori K, Kunugi H. Moderating effect of schizotypy on the relationship between smoking and neurocognition. *Eur Psychiatry.* 2013; 28: 457-462.
 86. Hori H*, Teraishi T, Sasayama D, Hattori K, Hashikura M, Higuchi T, Kunugi H. Relationship of temperament and character with cortisol reactivity to the combined dexamethasone/CRH test in depressed outpatients. *J Affect Disord.* 2013; 147: 128-136.
 87. Ota M*, Sato N, Ishikawa M, Hori H, Sasayama D, Hattori K, Teraishi T, Obu S, Nakata Y, Nemoto K, Moriguchi Y, Hashimoto R, Kunugi H. Discrimination of female schizophrenia patients from healthy women using multiple structural brain measures obtained with voxel-based morphometry. *Psychiatry Clin Neurosci.* 2012; 66: 611-617.
 88. Hori H*, Yamamoto N, Fujii T, Teraishi T, Sasayama D, Matsuo J, Kawamoto Y, Kinoshita Y, Ota M, Hattori K, Tatsumi M, Arima K, Kunugi H. Effects of the *CACNA1C* risk allele on neurocognition in patients with schizophrenia and healthy individuals. *Sci Rep.* 2012; 2: 634.
 89. Hori H, Kunugi H*. The efficacy of pramipexole, a dopamine receptor agonist, as an adjunctive treatment in treatment-resistant depression: an open-label trial. *ScientificWorldJournal.* 2012; 2012: 372474.
 90. Teraishi T, Ozeki Y, Hori H, Sasayama D, Chiba S, Yamamoto N, Tanaka H, Iijima Y, Matsuo J, Kawamoto Y, Kinoshita Y, Hattori K, Ota M, Kajiwara M, Terada S, Higuchi T, Kunugi H*. ¹³C-phenylalanine breath test detects altered phenylalanine kinetics in schizophrenia patients. *Transl Psychiatry.* 2012; 2: e119.
 91. Sasayama D*, Hattori K, Teraishi T, Hori H, Ota M, Yoshida S, Arima K, Higuchi T, Amano N, Kunugi H. Negative correlation between cerebrospinal fluid oxytocin levels and negative symptoms of male patients with schizophrenia. *Schizophr Res.* 2012; 139: 201-206.
 92. Hori H*, Teraishi T, Sasayama D, Hattori K, Ota M, Matsuo J, Kawamoto Y, Kinoshita Y, Kunugi H. Schizotypal trait in healthy women is associated with a shift away from dextrality on a spatial motor control task, but not on a force control task. *Psychiatry Res.* 2012; 200: 629-634.
 93. Hori H*, Matsuo J, Teraishi T, Sasayama D, Kawamoto Y, Kinoshita Y, Hattori K, Hashikura M, Higuchi T, Kunugi H. Schizotypy and genetic loading for schizophrenia impact upon neuropsychological status in bipolar II and unipolar major depressive disorders. *J Affect Disord.* 2012; 142: 225-232.

94. Sasayama D*, Hori H, Teraishi T, Hattori K, Ota M, Tatsumi M, Higuchi T, Amano N, Kunugi H. Possible impact of ADRB3 Trp64Arg polymorphism on BMI in patients with schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry*. 2012; 38: 341-344.
95. Okayasu H, Ozeki Y*, Fujii K, Okuri Y, Saeki Y, Hori H, Horie M, Higuchi T, Kunugi H, Shimoda K. Pharmacotherapeutic determinants for QTc interval prolongation in Japanese patients with mood disorder. *Pharmacopsychiatry*. 2012; 45: 279-283.
96. Ota M*, Ishikawa M, Sato N, Hori H, Sasayama D, Hattori K, Teraishi T, Nakata Y, Kunugi H. Glutamatergic changes in the cerebral white matter associated with schizophrenic exacerbation. *Acta Psychiatr Scand*. 2012; 126: 72-78.
97. Fujii T, Ota M, Hori H, Sasayama D, Hattori K, Teraishi T, Yamamoto N, Hashikura M, Tatsumi M, Higuchi T, Kunugi H*. Association between the functional polymorphism (C3435T) of the gene encoding P-glycoprotein (*ABCB1*) and major depressive disorder in the Japanese population. *J Psychiatr Res*. 2012; 46: 555-559.
98. Sasayama D*, Hori H, Teraishi T, Hattori K, Ota M, Matsuo J, Kawamoto Y, Kinoshita Y, Hashikura M, Amano N, Higuchi T, Kunugi H. More severe impairment of manual dexterity in bipolar disorder compared to unipolar major depression. *J Affect Disord*. 2012; 136: 1047-1052.
99. Hori H*, Teraishi T, Sasayama D, Fujii T, Hattori K, Ishikawa M, Kunugi H. Elevated cortisol level and cortisol/DHEAS ratio in schizophrenia as revealed by low-dose dexamethasone suppression test. *Open Neuropsychopharmacology Journal*. 2012; 5: 18-24.
100. Hattori K*, Tanaka H, Yamamoto N, Teraishi T, Hori H, Kinoshita Y, Matsuo J, Kawamoto Y, Kunugi H. Blood *CADPS2ΔExon3* expression is associated with intelligence and memory in healthy adults. *Biol Psychol*. 2012; 89: 117-122.
101. Hori H*, Teraishi T, Sasayama D, Matsuo J, Kawamoto Y, Kinoshita Y, Kunugi H. Relationships between season of birth, schizotypy, temperament, character and neurocognition in a non-clinical population. *Psychiatry Res*. 2012; 195: 69-75.
102. Sasayama D*, Hori H, Teraishi T, Hattori K, Ota M, Matsuo J, Kawamoto Y, Kinoshita Y, Amano N, Kunugi H. Association of cognitive performance with interleukin-6 receptor Asp358Ala polymorphism in healthy adults. *J Neural Transm*. 2012; 119: 313-318.
103. Fujii T, Yamamoto N, Hori H, Hattori K, Sasayama D, Teraishi T, Hashikura M, Tatsumi M, Okamoto N, Higuchi T, Kunugi H*. Support for association between the Ser205Leu polymorphism of *p75^{NTR}* and major depressive disorder. *J Hum Genet*. 2011; 56: 806-809.
104. Sasayama D*, Hori H, Teraishi T, Hattori K, Ota M, Iijima Y, Tatsumi M, Higuchi T, Amano N, Kunugi H. Possible association between *Interleukin-1beta* gene and schizophrenia in a Japanese population. *Behav Brain Funct*. 2011; 7: 35.
105. Sasayama D*, Hori H, Iijima Y, Teraishi T, Hattori K, Ota M, Fujii T, Higuchi T, Amano N, Kunugi H. Modulation of cortisol responses to the DEX/CRH test by polymorphisms of the interleukin-1beta gene in healthy adults. *Behav Brain Funct*. 2011; 7: 23.
106. Sasayama D, Wakabayashi C, Hori H, Teraishi T, Hattori K, Ota M, Ishikawa M, Arima K, Higuchi T, Amano N, Kunugi H*. Association of plasma IL-6 and soluble IL-6 receptor levels with the Asp358Ala polymorphism of the IL-6 receptor gene in schizophrenic patients. *J Psychiatr Res*. 2011; 45: 1439-1444.
107. Sasayama D*, Hori H, Teraishi T, Hattori K, Ota M, Matsuo J, Kawamoto Y, Kinoshita Y, Higuchi T, Amano N, Kunugi H. Association of interleukin-1β genetic polymorphisms with cognitive performance in elderly females without dementia. *J Hum Genet*. 2011; 56: 613-616.

108. Hattori K*, Tanaka H, Wakabayashi C, Yamamoto N, Uchiyama H, Teraishi T, Hori H, Arima K, Kunugi H. Expression of Ca²⁺-dependent activator protein for secretion 2 is increased in the brains of schizophrenic patients. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011; 35: 1738-1743.
109. Hori H, Teraishi T, Sasayama D, Ozeki Y, Matsuo J, Kawamoto Y, Kinoshita Y, Hattori K, Higuchi T, Kunugi H*. Poor sleep is associated with exaggerated cortisol response to the combined dexamethasone/CRH test in a non-clinical population. *J Psychiatr Res*. 2011; 45: 1257-1263.
110. Sasayama D*, Hori H, Teraishi T, Hattori K, Ota M, Matsuo J, Kawamoto Y, Kinoshita Y, Hashikura M, Koga N, Okamoto N, Sakamoto K, Higuchi T, Amano N, Kunugi H. Difference in Temperament and Character Inventory scores between depressed patients with bipolar II and unipolar major depressive disorders. *J Affect Disord*. 2011; 132: 319-324.
111. Hori H*, Teraishi T, Ozeki Y, Hattori K, Sasayama D, Matsuo J, Kawamoto Y, Kinoshita Y, Higuchi T, Kunugi H. Schizotypal personality in healthy adults is related to blunted cortisol responses to the combined dexamethasone/corticotropin-releasing hormone test. *Neuropsychobiology*. 2011; 63: 232-241.
112. Hori H*, Richards M, Kawamoto Y, Kunugi H. Attitudes toward schizophrenia in the general population, psychiatric staff, physicians, and psychiatrists: a web-based survey in Japan. *Psychiatry Res*. 2011; 186: 183-189.
113. Fujii T, Uchiyama H, Yamamoto N, Hori H, Tatsumi M, Ishikawa M, Arima K, Higuchi T, Kunugi H*. Possible association of the semaphorin 3D gene (*SEMA3D*) with schizophrenia. *J Psychiatr Res*. 2011; 45: 47-53.
114. Hori H, Ozeki Y, Teraishi T, Matsuo J, Kawamoto Y, Kinoshita Y, Suto S, Terada S, Higuchi T, Kunugi H*. Relationships between psychological distress, coping styles, and HPA axis reactivity in healthy adults. *J Psychiatr Res*. 2010; 44: 865-873.
115. Hashimoto R*, Hashimoto H, Shintani N, Ohi K, Hori H, Saitoh O, Kosuga A, Tatsumi M, Iwata N, Ozaki N, Kamijima K, Baba A, Takeda M, Kunugi H. Possible association between the pituitary adenylate cyclase-activating polypeptide (PACAP) gene and major depressive disorder. *Neurosci Lett*. 2010; 468: 300-302.
116. Hashimoto R*, Noguchi H, Hori H, Nakabayashi T, Suzuki T, Iwata N, Ozaki N, Kosuga A, Tatsumi M, Kamijima K, Harada S, Takeda M, Saitoh O, Kunugi H. A genetic variation in the dysbindin gene (*DTNBPI*) is associated with memory performance in healthy controls. *World J Biol Psychiatry*. 2010; 11: 431-438.
117. Ohi K, Hashimoto R*, Yasuda Y, Yamamori H, Hori H, Saitoh O, Tatsumi M, Takeda M, Iwata N, Ozaki N, Kamijima K, Kunugi H. No association between the *Bcl2-interacting killer (BIK)* gene and schizophrenia. *Neurosci Lett*. 2009; 463: 60-63.
118. Hashimoto R*, Noguchi H, Hori H, Ohi K, Yasuda Y, Takeda M, Kunugi H. The association analysis between the dysbindin gene (*DTNBPI*) and cognitive functions in Japanese subjects. *Psychiatry Clin Neurosci*. 2009; 63: 550-556.
119. Hashimoto R*, Ohi K, Okada T, Yasuda Y, Yamamori H, Hori H, Hikita T, Taya S, Saitoh O, Kosuga A, Tatsumi M, Kamijima K, Kaibuchi K, Takeda M, Kunugi H. Association analysis between schizophrenia and the AP-3 complex genes. *Neurosci Res*. 2009; 65: 113-115.
120. Hashimoto R*, Mori T, Nemoto K, Moriguchi Y, Noguchi H, Nakabayashi T, Hori H, Harada S, Kunugi H, Saitoh O, Ohnishi T. Abnormal microstructures of the basal ganglia in schizophrenia revealed by diffusion tensor imaging. *World J Biol Psychiatry*. 2009; 10: 65-69.
121. Hori H*, Ozeki Y, Terada S, Kunugi H. Functional near-infrared spectroscopy reveals altered hemispheric

- laterality in relation to schizotypy during verbal fluency task. *Prog Neuropsychopharmacol Biol Psychiatry*. 2008; 32: 1944-1951.
122. Izumi A, Iijima Y, Noguchi H, Numakawa T, Okada T, Hori H, Kato T, Tatsumi M, Kosuga A, Kamijima K, Asada T, Arima K, Saitoh O, Shiosaka S, Kunugi H*. Genetic variations of human Neuropeptide Y gene and psychiatric disorders: Polymorphism screening and possible association with bipolar disorder and cognitive functions. *Neuropsychopharmacology*. 2008; 33: 3237-3245.
 123. Richards M, Iijima Y, Shizuno T, Kamegaya Y, Hori H, Omori M, Arima K, Saitoh O, Kunugi H*. Failure to confirm an association between *Epsin 4* and schizophrenia in a Japanese population. *J Neural Transm*. 2008; 115: 1347-1354.
 124. Hashimoto R*, Moriguchi Y, Yamashita F, Mori T, Nemoto K, Okada T, Hori H, Noguchi H, Kunugi H, Ohnishi T. Dose-dependent effect of the Val66Met polymorphism of the brain-derived neurotrophic factor gene on memory-related hippocampal activity. *Neurosci Res*. 2008; 61: 360-367.
 125. Hori H*, Nagamine M, Soshi T, Okabe S, Kim Y, Kunugi H. Schizotypal traits in healthy women predict prefrontal activation patterns during a verbal fluency task: A near-infrared spectroscopy study. *Neuropsychobiology*. 2008; 57: 61-69.
 126. Noguchi H, Hori H*, Kunugi H. Schizotypal traits and cognitive function in healthy adults. *Psychiatry Res*. 2008; 161: 162-169.
 127. Hori H, Noguchi H, Hashimoto R, Nakabayashi T, Saitoh O, Murray RM, Okabe S, Kunugi H*. Personality in schizophrenia assessed with the Temperament and Character Inventory (TCI). *Psychiatry Res*. 2008; 160: 175-183.
 128. Hori H*, Noguchi H, Hashimoto R, Okabe S, Saitoh O, Kunugi H. IQ decline and memory impairment in Japanese patients with chronic schizophrenia. *Psychiatry Res*. 2008; 158: 251-255.
 129. Hashimoto R*, Noguchi H, Hori H, Ohi K, Yasuda Y, Takeda M, Kunugi H. A possible association between the Val158Met polymorphism of the catechol-O-methyl transferase gene and the personality trait of harm avoidance in Japanese healthy subjects. *Neurosci Lett*. 2007; 428: 17-20.
 130. Kunugi H*, Tanaka M, Hori H, Hashimoto R, Saitoh O, Hironaka N. Prepulse inhibition of acoustic startle response in Japanese patients with chronic schizophrenia. *Neurosci Res*. 2007; 59: 23-28.
 131. Hashimoto R*, Hashimoto H, Shintani N, Chiba S, Hattori S, Okada T, Nakajima M, Tanaka K, Kawagishi N, Nemoto K, Mori T, Ohnishi T, Noguchi H, Hori H, Suzuki T, Iwata N, Ozaki N, Nakabayashi T, Saitoh O, Kosuga A, Tatsumi M, Kamijima K, Weinberger DR, Kunugi H, Baba A. Pituitary adenylate cyclase-activating polypeptide is associated with schizophrenia. *Mol Psychiatry*. 2007; 12: 1026-1032.
 132. Fujii T, Iijima Y, Kondo H, Shizuno T, Hori H, Nakabayashi T, Arima K, Saitoh O, Kunugi H*. Failure to confirm an association between the *PLXNA2* gene and schizophrenia in a Japanese population. *Prog Neuropsychopharmacol Biol Psychiatry*. 2007; 31: 873-877.
 133. Mori T, Ohnishi T*, Hashimoto R, Nemoto K, Moriguchi Y, Noguchi H, Nakabayashi T, Hori H, Harada S, Saitoh O, Matsuda H, Kunugi H. Progressive changes of white matter integrity in schizophrenia revealed by diffusion tensor imaging. *Psychiatry Res*. 2007; 154: 133-145.
 134. Richards M, Iijima Y, Kondo H, Shizuno T, Hori H, Arima K, Saitoh O, Kunugi H*. Association study of the vesicular monoamine transporter 1 (VMAT1) gene with schizophrenia in a Japanese population. *Behav Brain Funct*. 2006; 2: 39.
 135. Kunugi H*, Hashimoto R, Okada T, Hori H, Nakabayashi T, Baba A, Kudo K, Omori M, Takahashi S,

- Tsukue R, Anami K, Hirabayashi N, Kosuga A, Tatsumi M, Kamijima K, Asada T, Harada S, Arima K, Saitoh O. Possible association between nonsynonymous polymorphisms of the anaplastic lymphoma kinase (ALK) gene and schizophrenia in a Japanese population. *J Neural Transm*. 2006; 113: 1569-1573.
136. Hori H*, Noguchi H, Hashimoto R, Nakabayashi T, Omori M, Takahashi S, Tsukue R, Anami K, Hirabayashi N, Harada S, Saitoh O, Iwase M, Kajimoto O, Takeda M, Okabe S, Kunugi H. Antipsychotic medication and cognitive function in schizophrenia. *Schizophr Res*. 2006; 86: 138-146.
137. Ohnishi T*, Hashimoto R, Mori T, Nemoto K, Moriguchi Y, Iida H, Noguchi H, Nakabayashi T, Hori H, Ohmori M, Tsukue R, Anami K, Hirabayashi N, Harada S, Arima K, Saitoh O, Kunugi H. The association between the Val158Met polymorphism of the catechol-O-methyl transferase gene and morphological abnormalities of the brain in chronic schizophrenia. *Brain*. 2006; 129: 399-410.

和文原著論文 (直近のものから順に記載)

1. 太田深秀, 野田隆政, 佐藤典子, 服部功太郎, 堀 弘明, 篠山大明, 寺石俊也, 長島杏奈, 大部聡子, 樋口輝彦, 功刀 浩. メランコリー型の特徴を伴う大うつ病性障害と非定型の特徴を伴う大うつ病性障害の白質障害の違い. *精神神経学雑誌* 2017; 119: 295-302.
2. 太田深秀, 若林千里, 松尾淳子, 木下 裕紀子, 堀 弘明, 服部 功太郎, 篠山大明, 寺石俊也, 大部聡子, 小澤隼人, 大久保 勉, 功刀 浩. 健常者の感覚運動フィルター機構に対する L-テアニンの影響. *精神神経学雑誌* 2015; 117: 813-819.
3. 太田深秀, 佐藤典子, 石川正憲, 堀 弘明, 篠山大明, 服部 功太郎, 寺石俊也, 大部聡子, 中田安弘, 根本清貴, 守口善也, 橋本亮太, 功刀 浩. MRIによる女性統合失調症患者と女性健常群との判別分析. *精神神経学雑誌* 2013; 115: 1171-1177.
4. 古賀賀恵, 服部 功太郎, 堀 弘明, 功刀 浩. 緑茶、コーヒーを飲む習慣と大うつ病との関連. *New Diet Therapy* 2013; 29: 31-38.
5. 功刀 浩, 柳沢洋美, 田中美穂, 岡本洋平, 堀 弘明, 橋本亮太, 中林哲夫, 岡本長久, 大森まゆ, 沢村香苗, 斎藤 治, 樋口輝彦, 廣中直行. 統合失調症とうつ病におけるプレパルスインビジョンと関連指標:予備的報告. *精神医学* 2007; 49: 253-260.
6. 野口広子, 堀 弘明, 橋本亮太, 中林哲夫, 津久江 亮太郎, 穴見公孝, 斎藤 治, 功刀 浩. 統合失調症における知的機能の縦断的变化について:予備的検討. *Progress in Medicine* 2006; 26: 1709-1711.
7. 今岡岳史, 堀 達, 小宮山 徳太郎, 村松玲美, 堀 弘明, 高木希奈. 退院後抗酒剤服薬支援を要した患者の特徴と予後. *日本アルコール精神医学雑誌* 2005; 11: 25-29.

[総説論文]

英文総説論文 (直近のものから順に記載. *: corresponding author.)

1. Hakamata Y*, Suzuki Y, Kobashikawa H, Hori H*. Neurobiology of early life adversity: A systematic review of meta-analyses towards an integrative account of its neurobiological trajectories to mental disorders. *Front Neuroendocrinol*. 2022; 65: 100994.
2. Hori H*, Kim Y. Inflammation and post-traumatic stress disorder. *Psychiatry Clin Neurosci*. 2019; 73: 143-153. (招待総説『PCN Frontier Review』) <“Highly Cited Paper” in Web of Science>
3. Kunugi H*, Hori H, Ogawa S. Biochemical markers subtyping major depressive disorder. *Psychiatry*

Clin Neurosci. 2015; 69: 597-608.

4. Hori H, Kunugi H*. Dopamine agonist-responsive depression. *Psychogeriatrics.* 2013; 13: 189-195.
5. Abbs B, Achalia RM, Adelufosi AO, Aktener AY, Beveridge NJ, Bhakta SG, Blackman RK, Bora E, Byun MS, Cabanis M, Carrion R, Castellani CA, Chow TJ, Dmitrzak-Weglarz M, Gayer-Anderson C, Gomes FV, Haut K, Hori H, Kantrowitz JT, Kishimoto T, Lee FH, Lin A, Palaniyappan L, Quan M, Rubio MD, Ruiz de Azúa S, Sahoo S, Strauss GP, Szczepankiewicz A, Thompson AD, Trotta A, Tully LM, Uchida H, Velthorst E, Young JW, O'Shea A, Delisi LE*. The 3rd Schizophrenia International Research Society Conference, 14-18 April 2012, Florence, Italy: summaries of oral sessions. *Schizophr Res.* 2012; 141: e1-e24.
6. Numakawa T*, Yokomaku D, Richards M, Hori H, Adachi N, Kunugi H. Functional interactions between steroid hormones and neurotrophin BDNF. *World J Biol Chem.* 2010; 1: 133-143.
7. Kunugi H*, Hori H, Adachi N, Numakawa T. Interface between hypothalamic-pituitary-adrenal axis and brain-derived neurotrophic factor in depression. *Psychiatry Clin Neurosci.* 2010; 64: 447-459.
8. Baharnoori M, Bartholomeusz C, Boucher AA, Buchy L, Chaddock C, Chiliza B, Föcking M, Fornito A, Gallego JA, Hori H, Huf G, Jabbar GA, Kang SH, El Kissi Y, Merchán-Naranjo J, Modinos G, Abdel-Fadeel NA, Neubeck AK, Ng HP, Novak G, Owolabi OO, Prata DP, Rao NP, Riecanaky I, Smith DC, Souza RP, Thienel R, Trotman HD, Uchida H, Woodberry KA, O'Shea A, DeLisi LE*. The 2nd Schizophrenia International Research Society Conference, 10-14 April 2010, Florence, Italy: summaries of oral sessions. *Schizophr Res.* 2010; 124: e1-62.
9. Hashimoto R*, Hattori S, Chiba S, Yagasaki Y, Okada T, Kumamaru E, Mori T, Nemoto K, Tani H, Hori H, Noguchi H, Numakawa T, Ohnishi T, Kunugi H. Susceptibility genes for schizophrenia. *Psychiatry Clin Neurosci.* 2006; 60 suppl 1: S4-S10.

和文総説論文 (直近のものから順に記載)

1. 堀 弘明: 心的外傷後ストレス症における栄養学的問題. *精神医学* 2024; 66: 295-301.
2. 堀 弘明: 複雑性 PTSD の神経生物学 -逆境的小児期体験(ACEs)と心的外傷後ストレス症(PTSD)の研究からの考察-. *精神医学* 2023; 65: 1172-1182.
3. 堀 弘明: PTSD への薬物療法. *精神科* 2023; 43: 209-217.
4. 堀 弘明: うつ病の発症におけるストレスの役割. *日本生物学的精神医学会誌* 2022; 33: 148-154.
5. 堀 弘明: PTSD の統合的理解を目指した心理学的・生物学的研究. *トラウマティック・ストレス* 2022; 20: 11-19.
6. 堀 弘明, 金 吉晴: 認知と炎症に着目した PTSD の遺伝学的検討. *日本生物学的精神医学会誌* 2022; 33: 10-15.
7. 堀 弘明: 複雑性 PTSD の神経生物学. *精神療法* 2021; 47: 463-472.
8. 堀 弘明: PTSD と炎症. *総合病院精神医学* 2019; 31: 160-173.
9. 堀 弘明: PTSD 治療における薬物療法の位置づけ. *臨床精神薬理* 2017; 20: 789-796.
10. 堀 弘明: PTSD の遺伝研究. *トラウマティック・ストレス* 2015; 13: 150-159.
11. 堀 弘明, 功刀 浩: 初発精神病エピソードの初回面接. *精神科治療学* 2014; 29: 911-917.

12. 堀 弘明, 寺石俊也, 篠山大明, 服部功太郎, 橋倉 都, 樋口輝彦, 功刀 浩: 大うつ病性障害におけるパーソナリティはコルチゾール反応性を予測する:デキサメタゾン/CRH テストによる検討. 日本神経精神薬理学雑誌 2013; 33: 61-63.
13. 功刀 浩, 堀 弘明: うつ病のバイオマーカーと亜型 特に視床下部-下垂体-副腎系機能との関連について. 臨床精神医学 2013; 42: 991-998.
14. 功刀 浩, 堀 弘明, 沼川忠広, 太田深秀: 視床下部-下垂体-副腎系とうつ病 最近の展開. 日本神経精神薬理学雑誌 2012; 32: 203-209.
15. 堀 弘明, 功刀 浩: ストレス反応・うつ病とニューロペプチド. HORMONE FRONTIER IN GYNECOLOGY 2010; 17: 31-37.
16. 堀 弘明, 功刀 浩: 視床下部-下垂体-副腎系とうつ病 : 最近の知見. 分子精神医学 2008; 8: 17-23.
17. 堀 弘明, 功刀 浩: 統合失調症の神経生理学的研究. メディカル・サイエンス・ダイジェスト 2008; 34: 582-585.

[著書]

分担執筆著書 (直近のものから順に記載)

1. Hori H, Hakamata Y. “Chapter 8. Inflammation and traumatic stress.” Stress: Immunology and Inflammation: Handbook of Stress Series Volume 5, Fink G (Editor), Academic Press, London, pp 65-75, 2023.
2. 堀 弘明:「第8章 第2節 心的外傷後ストレス障害」. 精神保健医療福祉白書編集委員会(編): 精神保健医療福祉白書 2018/2019. 中央法規出版, 東京, p159, 2018.
3. 堀 弘明:「第8章 第2節 心的外傷後ストレス障害」. 精神保健医療福祉白書編集委員会(編): 精神保健医療福祉白書 2017. 中央法規出版, 東京, p174, 2016.
4. 堀 弘明, 功刀 浩:「ミルタザピンのバイオマーカーに対する影響」. ミルタザピンのすべて, 編集: 小山司, 樋口輝彦. 先端医学社, 東京, pp 176-183, 2012.
5. 堀 弘明, 功刀 浩:「視床下部-下垂体-副腎系とうつ病スペクトラム」. 気分障害の薬理・生化学, 編集: 躁うつ病の薬理・生化学的研究懇話会. 医薬ジャーナル社, 東京, pp 174-191, 2012.
6. 堀 弘明, 功刀 浩:「視床下部-下垂体-副腎系機能検査法と精神疾患」. 精神疾患診断のための脳形態・機能検査法, 編集: 三國雅彦, 福田正人, 功刀浩. 新興医学出版社, 東京, pp 91-100, 2012.
7. 堀 弘明, 功刀 浩:「統合失調症型パーソナリティと半球優位性」. 精神疾患とNIRS, 編集: 福田正人. 中山書店, 東京, pp 202-211, 2009.
8. 堀 弘明, 功刀 浩:「第8章 V. 神経内分泌学」. 気分障害, 編集: 上島国利, 樋口輝彦, 野村 総一郎, 大野 裕, 神庭重信, 尾崎紀夫. 医学書院, 東京, pp 245-252, 2008.

翻訳書 (直近のものから順に記載)

1. 堀 弘明 (訳) : モデルで考える精神疾患. (Peter Tyrer, Derek Steinberg: Models for Mental Disorder), 星和書店, 東京, 2012.
2. 功刀 浩, 堀 弘明 (共訳) : 統合失調症の常識は本当か? (Colm McDonald, Katja Schulze, Robin Murray, Pdraig Wright: Schizophrenia: Challenging the Orthodox), 培風館, 東京, 2009.
3. 功刀 浩, 堀 弘明 (共訳) : 統合失調症 100 の Q&A. (Lynn E. Delisi: 100 Questions & Answers About Schizophrenia: Painful Minds), 星和書店, 東京, 2008.

[国際学会発表] (直近のものから順に、筆頭演者分のみ記載)

招待講演・シンポジウム

1. Hori H, Kim Y: A genetic study of PTSD focusing on cognition and inflammation. ISTSS 37th Annual Meeting, Online • Live, 2021.11.2-5. (シンポジウム)
2. Hori H: Cognitive dysfunction in schizophrenia and other psychiatric disorders: potential role of inflammation. WFSBP2018KOBE, Kobe, 2018.9.7-9. (シンポジウム)
3. Hori H: Hypothalamic-pituitary-adrenal axis function and gene expression profiles as useful blood biomarkers for major depressive disorders. The 10th Annual Scientific Meeting of Hong Kong Society of Biological Psychiatry, Hong Kong, 2017.3.11. (日本生物学的精神医学会 2016 年度若手国際交流プログラムによる派遣・招待講演)
4. Hori H: A multifaceted approach to understanding the pathophysiology of schizophrenia from the perspective of schizotypal personality. The 2012 Annual Meeting of Taiwanese Society of Biological Psychiatry and Neuropsychopharmacology, Taipei, Taiwan, 2012.9.22. (日本生物学的精神医学会 2012 年度若手国際交流プログラムによる派遣・招待講演)

一般演題

1. Hori H, Itoh M, Lin M, Niwa M, Ino K, Imai R, Ogawa S, Sekiguchi A, Kunugi H, Kamo T, Kim H: Inflammatory markers in adult women with post-traumatic stress disorder (PTSD). 13th World Congress of Biological Psychiatry, Copenhagen, 2017.6.18-22. (ポスター)
2. Hori H, Sasayama D, Teraishi T, Yamamoto N, Nakamura S, Ota M, Hattori K, Kim Y, Higuchi T, Kunugi H: Involvement of hypothalamic-pituitary-adrenal axis in the mechanism of transcriptome-wide identified differentially expressed genes for depression. 29th ECNP Congress, Vienna, 2016.9.17-20. (ポスター)
3. Hori H, Sasayama D, Teraishi T, Yamamoto N, Nakamura S, Ota M, Hattori K, Kim Y, Higuchi T, Kunugi H: Blood-based gene expression signatures of medication-free outpatients with major depressive disorder: integrative genome-wide and candidate gene analyses identify genes suggestive of molecular mechanisms underlying depression. World Psychiatric Association International Congress 2015, Taipei, 2015.11.18-20. (ポスター)
4. Hori H, Yamamoto N, Fujii T, Teraishi T, Sasayama D, Arima K, Kunugi H: The bipolar risk allele at CACNA1C impacts on neurocognition in patients with schizophrenia. The 2nd Meeting of East Asian Bipolar Forum, Fukuoka, 2012.9.7-8. (口演)

5. Hori H, Teraishi T, Sasayama D, Hattori K, Hashikura M, Higuchi T, Kunugi H: Personality dimensions in major depressive disorder predict cortisol reactivity to the combined dexamethasone/CRH test. The 28th CINP World Congress of Neuropsychopharmacology, Stockholm, Sweden, 2012.6.3-7. (ポスター)
6. Hori H, Matsuo J, Teraishi T, Sasayama D, Kawamoto Y, Kinoshita Y, Hattori K, Hashikura M, Higuchi T, Kunugi H: Schizotypy and genetic loading for schizophrenia impact upon neuropsychological status in bipolar II and unipolar major depressive disorders. Society of Biological Psychiatry 67th Annual Scientific Convention, Philadelphia, USA, 2012.5.3-5. (ポスター)
7. Hori H, Matsuo J, Teraishi T, Sasayama D, Kawamoto Y, Kinoshita Y, Hattori K, Hashikura M, Higuchi T, Kunugi H: Schizotypy and genetic loading for schizophrenia impact upon neuropsychological status in bipolar II and unipolar major depressive disorders. The 3rd Biennial Schizophrenia International Research Conference, Florence, Italy, 2012.4.14-18. (ポスター)
8. Hori H, Teraishi T, Sasayama D, Ishikawa M, Higuchi T, Kunugi H: Elevated cortisol levels and cortisol/dehydroepiandrosterone ratio in schizophrenia as revealed by the dexamethasone suppression test. 13th International Congress on Schizophrenia Research, Colorado, USA, 2011.4.2-6. (ポスター)
9. Hori H, Teraishi T, Ozeki Y, Matsuo J, Kawamoto Y, Kinoshita Y, Suto S, Higuchi T, Kunugi H: Schizotypal personality in healthy adults is related to blunted HPA axis reactivity. The 2nd Biennial Schizophrenia International Research Conference, Florence, Italy, 2010.4.10-14. (ポスター)
10. Hori H, Richards M, Kunugi H: Attitudes towards schizophrenia in the general population, psychiatric staff, physicians, and psychiatrists: A web-based survey in Japan. The 15th Biennial Winter Workshop in Psychoses, Barcelona, Spain, 2009.11.15-18. (ポスター)
11. Hori H, Ozeki Y, Higuchi T, Kunugi H: Relationships of psychological distress and sleep status with HPA axis function in a non-clinical population. World Psychiatric Association, Florence, Italy, 2009.4.1-4. (ポスター)
12. Hori H, Noguchi H, Hashimoto R, Nakabayashi T, Saitoh O, Murray RM, Okabe S, Kunugi H: Personality traits assessed with the Temperament and Character Inventory in schizophrenia. International Congress on Schizophrenia Research, Colorado, USA, 2007.3.28-4.1. (ポスター)

[国内学会発表 (シンポジウム・ワークショップ)] (直近のものから順に、筆頭演者分のみ記載)

1. 堀 弘明, 金 吉晴: メマンチンによる PTSD 治療の試み. シンポジウム S-3, PTSD に対する薬物療法: 今後の展望を考える. 第 21 回日本トラウマティック・ストレス学会, オンライン, 2022.7.23-24. (シンポジウム)
2. 堀 弘明, 金 吉晴: 遺伝環境相互作用に着目した PTSD の病因理解. シンポジウム 6, 恐怖記憶の分子・生理学的基盤の解明と PTSD の治療開発. 第 118 回日本精神神経学会学術総会, 福岡/オンライン, 2022.6.16-18. (シンポジウム)
3. 堀 弘明: PTSD の統合的理解を目指した心理学的・生物学的研究. シンポジウム S-14, PTSD 病態の理解と新規治療方法の開発. 第 20 回日本トラウマティック・ストレス学会, オンライン・オンデマンド配信, 2021.7.17-18. (シンポジウム)
4. 堀 弘明, 金 吉晴: 認知と炎症に着目した PTSD の遺伝学的検討. シンポジウム 19, PTSD 病態の理解と新規治療方法の開発. 第 43 回日本生物学的精神医学会/第 51 回日本神経精神薬理学会合同年会, 京都/オンライン, 2021.7.14-16. (シンポジウム)

5. 堀 弘明：うつ病の発症におけるストレスの役割. シンポジウム 8, うつ病の解明と新規治療標的の探索～基礎と臨床の双方向アプローチ～. 第 43 回日本生物学的精神医学会／第 51 回日本神経精神薬理学会合同年会, 京都／オンライン, 2021.7.14-16. (シンポジウム)
6. 堀 弘明: A multifaceted approach to understanding the pathogenesis of PTSD: from genes, biomarkers to phenotypes. 第 60 回日本神経化学学会大会, 仙台, 2017.9.7-9. (シンポジウム)
7. 堀 弘明：うつ病バイオマーカーとしての視床下部-下垂体-副腎系. 第 13 回日本うつ病学会総会, 愛知, 2016.8.5-6. (シンポジウム)
8. 堀 弘明, 功刀 浩：外来うつ病、健常成人における DEX/CRH テスト. 日本心理学会第 71 回大会, 東京, 2007.9.18-20. (ワークショップ)

[研修会講師] (直近のものから順に記載)

1. 堀 弘明：PTSD の神経科学と薬物療法. 令和 5 年度 PTSD 対策専門研修 B.専門コース 2, オンライン, 2024.1.17-18.
2. 堀 弘明：PTSD の神経科学と薬物療法. 令和 5 年度 PTSD 対策専門研修 B.専門コース 1, オンライン, 2023.12.20-21.
3. 堀 弘明：PTSD の神経科学と薬物療法. 令和 4 年度 PTSD 対策専門研修 B.専門コース 2, オンライン, 2023.1.19-20.
4. 堀 弘明：PTSD の神経科学と薬物療法. 令和 4 年度 PTSD 対策専門研修 B.専門コース 1, オンライン, 2022.12.15-16.
5. 堀 弘明：PTSD の神経科学と薬物療法. 令和 3 年度 PTSD 対策専門研修 B.専門コース 2, オンライン, 2022.1.13-14.
6. 堀 弘明：PTSD の神経科学と薬物療法. 令和 3 年度 PTSD 対策専門研修 B.専門コース 1, オンライン, 2021.12.16-17.
7. 堀 弘明：PTSD の神経科学と薬物療法. 令和 2 年度 PTSD 対策専門研修 B.専門コース 2, オンライン開催, 2021.1.21-22.
8. 堀 弘明：PTSD の神経科学と薬物療法. 令和 2 年度 PTSD 対策専門研修 B.専門コース 1, オンライン開催, 2020.12.3-4.
9. 堀 弘明：PTSD の神経科学と薬物療法. 令和元年度 PTSD 対策専門研修 B.専門コース 2, 東京, 2019.12.19-20.
10. 堀 弘明：PTSD の神経科学と薬物療法. 令和元年度 PTSD 対策専門研修 B.専門コース 1, 東京, 2019.11.28-29.
11. 堀 弘明：PTSD の神経科学と薬物療法. 平成 30 年度 PTSD 対策専門研修 B.専門コース 2. 東京: 2019.2.19
12. 堀 弘明：PTSD の神経科学と薬物療法. 平成 30 年度 PTSD 対策専門研修 B.専門コース 1. 東京: 2019.2.5
13. 堀 弘明：PTSD の神経科学と薬物療法. 平成 29 年度 PTSD 対策専門研修 応用コース A, 東京, 2018.1.23.
14. 堀 弘明：PTSD の生物学的基盤と薬物療法. 平成 28 年度 PTSD 対策専門研修事業 A.通常コ

ース, 東京, 2017.2.22.

15. 堀 弘明: PTSD の生物学的基盤と薬物療法. 平成 27 年度 PTSD 対策専門研修事業 A.通常コース, 東京, 2016.2.11.

[プレスリリース] (直近のものから順に記載)

1. 堀 弘明, 袴田優子: 小児期の虐待体験が免疫システムに関連することを発見 ~概日リズム平坦化がその鍵を握る可能性~|日本の研究.com. アンドラ, 2022.3.7.
2. 堀 弘明, 袴田優子: 小児期の被虐待体験、成人後の IL-6 濃度日内変動の平坦化に関連の可能性-NCNP ほか. QLifePro, 2022.3.3.
3. 金 吉晴, 堀 弘明: メマンチンによる PTSD 治療の有効性. ケアネット, 2021.5.4.
4. 金 吉晴, 堀 弘明, 伊藤真利子, 林明明: マイナビニュース. 幼少時の心理的虐待と成人後の注意バイアス変動性の関連性を NCNP などが確認, 2021.3.16.
5. 金 吉晴, 堀 弘明: ヨミドクター. 認知症治療薬メマンチンが PTSD に有望か 心理療法に匹敵する効果、臨床試験を計画へ, 2021.3.10.
6. 金 吉晴, 堀 弘明: Medical Tribune. 認知症治療薬メマンチンが PTSD に有望か, 2021.3.4.
7. 金 吉晴, 堀 弘明: 薬事ニュース(CB news). PTSD 治療, メマンチンの有効性明らかに NCNP 精神保健研究所の研究グループ, 2021.2.5.
8. 金 吉晴, 堀 弘明, 他: 毎日新聞, 南日本新聞, 山梨日日新聞, 福島民報, 他. 抗認知症薬、PTSD に効果か 体験の認識を改善 国立精神研チーム, 2021.1.25.
9. 金 吉晴, 堀 弘明: 日本経済新聞デジタル版. 国立精神・神経医療研究センター、心的外傷後ストレス障害の治療におけるメマンチンの有効性 -オープンラベル臨床試験による実証-, 2021.1.25.
10. 金 吉晴, 堀 弘明, 大塚豪士: QLifePro. PTSD の症状や認知機能に CRP 遺伝子多型が関与することを発見, 2020.12.15.
11. 金 吉晴, 堀 弘明, 中山未知: 科学新聞. 幼少期の性的虐待 成人女性に長期的悪影響, 2020.7.17.

[特許取得]

1. 中村誠二, 石澤洋平, 的場 亮, 松原謙一, 功刀 浩, 堀 弘明: 特許「気分障害を検出する方法」, 特許第 7084580 号, 2022 年 6 月 7 日登録, 日本

[雑誌コラム・監修等]

1. 堀 弘明: PTSD の分子生物学. 精神科 Resident 2022; 3: 39-40. (コラム)
2. 堀 弘明, 金 吉晴: 不安症と心的外傷後ストレス障害 (PTSD) の相違点: 何故、DSM-5 で PTSD は不安症から独立したのか? 精神科 Resident 2022; 3: 46-48. (コラム)
3. 堀 弘明: 幻覚剤を PTSD 治療に. 日経サイエンス 2022; 52: 64-70. (監修)
4. 堀 弘明: 幻覚剤を PTSD 治療に. 別冊日経サイエンス 2022; 256: 68-73. (監修)